



Elo Water Pte Ltd

Hyflux Innovation Centre, 80 Bendemeer Road, Singapore 339949
(65) 3157 8080 | www.elowatershop.com



HYPOXIA

AND
ITS EFFECTS ON HEALTH



WHY IS OXYGEN IMPORTANT?

Oxygen is important to every cell in your body. Without it, your cells cannot make much energy, and their metabolism is less effective.

Within every cell there are thousands of little power-plants called mitochondria^a. These are 'power plants' that generate energy in the form of ATP^b. In order to fire up the power-plants to make the energy molecule ATP, in a process called oxidative phosphorylation (OXPHOS)^c, you will need a generous supply of oxygen to the cells¹.

When you do not have enough oxygen, you switch from a highly efficient energy-making machine (OXPHOS) to being 16 times less efficient by using a non-oxygen energy process called anaerobic glycolysis^d and in doing so, you

generate a lot of lactic acid, the substance that makes your muscles feel sore and tired².

Athletes and sports enthusiasts are familiar with these processes that determine how well you perform and how well you recover from exercise. The more ATP you can create, the more energy your muscles get to use for exercise, translating into greater speed and strength of muscle contraction and longer endurance³. With more oxygen, you generate 16 times more energy using a process (OXPHOS) that does not produce lactic acid¹. In other words, you create more energy to power your exercise and you will end up with better post recovery with less muscle soreness.

Without sufficient oxygen, our cells have much less energy and perform their functions less efficiently⁴. This can be a problem because energy is needed to keep the cells healthy and optimally functioning. When cells are not functioning well, this is when the organs start to malfunction and bring on the start of disease. Cells that are starved of oxygen can also start to break down and die prematurely, bringing on degeneration⁵. Degeneration of cells drives the aging process in your body and leads to degenerative diseases such as dementia, retinal degeneration^{4,6}. When a person ages, degenerative processes cause reduction in lung capacity, hardening and blockage of the arteries, resulting in a less effective delivery system to bring oxygen to the body cells⁶. This creates a vicious cycle of aging and degeneration.

In conclusion, oxygen is an important universal nutrient that determines efficiency of energy supply to the living cells for powering energy intensive cell activities and maintaining overall wellbeing of cells.

Definitions

- a) **Mitochondria** – these are tiny components inside a living cell responsible for generating energy using oxygen. They are like power generators for the living cell, making sure that the cell has a constant energy supply in order to stay alive and carry out its normal functions.
- b) **ATP** – Adenosine triphosphate, an energy-carrying molecule found in the cells of all living things. ATP captures chemical energy obtained from the breakdown of food molecules and releases it to fuel all cellular processes needed for life. It is the source of energy that keeps everything going.
- c) **OXPHOS** - Oxidative Phosphorylation is process whereby a cell uses oxygen to generate large amounts of energy in the form of ATP. This process, which takes place in mitochondria, is the major source of ATP in aerobic organisms.
- d) **Anaerobic glycolysis** - Anaerobic glycolysis is a method used by cells to produce ATP by transforming glucose into lactate when limited amounts of oxygen (O₂) are available. Compared to OXPHOS, anaerobic glycolysis creates 16 times less energy.

References:

1. Oxidative Phosphorylation. D.A. Bender. Encyclopedia of Food Sciences and Nutrition (Second Edition), 2003
2. Biochemistry of exercise-induced metabolic acidosis. Am J Physiol Regul Integr Comp Physiol. 2004 Sep;287(3):R502-16.
3. Energy supply and muscle fatigue in humans. Acta Physiol Scand. 1998 Mar;162(3):261-6.
4. Boveris A Navarro A. Brain mitochondrial dysfunction in aging. IUBMB Life . 2008;60:308-314.
5. Navarro A Boveris A. The mitochondrial energy transduction system and the aging process. Am J Physiol Cell Physiol . 2007;292:C670-C686.
6. Physiology Considerations in the Geriatric Patient. Anesthesiol Clin. 2015 September ; 33(3): 447-456.

WHAT IS HYPOXIA?

Hypoxia is a medical term used to describe a condition where bodily tissues are not receiving enough oxygen. In other words, hypoxia literally means low oxygen levels in your body.

Hypoxia can be present in anyone because of a variety of reasons, and it is associated with many different medical conditions. It can be caused by decreased breathable oxygen in the air (like when you are travelling in an airplane or on a high mountain), decreased ability for the blood to carry oxygen to the organ tissues (such as in anemia, thalassemia^a, severe blood loss or blocked arteries or capillaries), decreased ability of tissues to absorb oxygen (like lung disease) or decreased ability of cells to utilize oxygen (such as carbon monoxide^b poisoning or abnormal mitochondria^c function).

When a person becomes older, reduction in lung capacity, hardening and blockage of their arteries and small vessels may compromise the ability to fully oxygenate the body tissues. Environmental pollution, smoking and suboptimal breathing patterns can all contribute to poorer oxygen levels in the body.

Oftentimes, hypoxia can exist in isolated organs or areas in the body affected by poor blood supply or abnormal behavior of the cells that have difficulty using oxygen. Many common diseases are associated with isolated hypoxia in affected organs. While generalized hypoxia can be measured by special oxygen measurement devices in the hospital, isolated hypoxia inside organs or within deep tissue can only be measured invasively and is therefore not commonly performed.

Definitions

a) **Thalassemia** – a hereditary, genetic blood disorder common in people of Mediterranean and Southeast Asian descent, where the oxygen-carrying red pigment inside red blood cells (hemoglobin) is structured abnormally and therefore unable to effectively carry oxygen.

b) **Carbon monoxide** is a gas present in car exhaust fumes and cigarette smoke. This gas binds to red blood cells and prevents the red cells from carrying oxygen.


c) **Mitochondria** – these are tiny components inside a living cell responsible for generating energy using oxygen. They are like power generators for the living cell, making sure that the cell has a constant energy supply in order to stay alive and carry out its normal functions.

THE CONNECTION BETWEEN HYPOXIA AND DIABETES

The lack of oxygen, or hypoxia, would appear very far removed from any connections with diabetes mellitus, a condition of abnormal glucose metabolism.

Indeed, in recent years, medical scientists are beginning to find out that hypoxia is closely associated with disease processes in diabetes, and is not only the result of abnormal blood circulation present in diabetes, but is also responsible for driving progression of some diabetic complications¹.

Fat cell hypoxia is clearly a factor that contributes to the development of insulin resistance and eventually diabetes. The role that hypoxia plays in the causation of a diabetic state was made clearer through the work of researchers at the University of California, San Diego School of Medicine. They discovered that eating fatty foods causes activation of a protein inside the fat cell membrane, called adenine nucleotide translocase 2 (ANT2), which consumes huge amounts of oxygen, leaving very little for the rest of the cell².



When a cell is oxygen starved (hypoxic) it becomes 'stressed' and will release nasty substances called 'chemokines'^a that turn on the immune system's inflammatory response^a. When this happens, the resulting low-grade tissue inflammation will impair the way cells interact with insulin, leading to insulin resistance, a state where insulin gets less and less effective in lowering blood sugar. This precedes the development of type 2 diabetes².

Diabetes patients suffer poor blood flow through the narrowed small blood vessels (microangiopathy^b) all over their body and this means that the oxygen carried in red blood cells have difficulty getting through these narrowed vessels to reach the body tissue.



In a clinical study conducted in the Netherlands, researchers found that people with diabetes, when compared to healthy subjects without diabetes, have consistently lower tissue oxygen values measured at various sites of the body³. They deduce that this could be due to the narrowed blood vessels supplying inadequate oxygen to the body cells, causing hypoxia.³ The doctors at the Hyperbaric Unit, Royal Adelaide Hospital, The University of Adelaide, South Australia also discovered that increasing the patient's body oxygen levels using hyperbaric oxygen therapy^c led to better response of the body to insulin with more effective blood sugar lowering, further drawing the link between diabetes and body oxygen levels⁴.

Obstructive sleep apnea (OSA) occurs when your throat muscles intermittently relax and block your airway during sleep, commonly observed as snoring. This causes breathing to repeatedly stop and start during sleep, leading to intermittent lack of oxygen to the brain. Severe obstructive sleep apnea (OSA) increases a person's risk of developing diabetes by 30% or more, according to a study of 11,000 patients, published in the American Journal of Respiratory and Critical Care Medicine⁵. Intermittent hypoxia in people with obstructive sleep apnea is found to propagate insulin resistance and increase the risks of developing diabetes⁶. Now that that link has been confirmed in such a large trial with a long follow-up period⁵, clinicians may be able to intervene and take diabetes prevention measures for patients with OSA who have not yet developed the disease.



References:

1. Mechanisms of Disease: the hypoxic tubular hypothesis of diabetic nephropathy. *Nature Clinical Practice Nephrology* volume 4, pages 216–226 (2008)
2. Increased Adipocyte O₂ Consumption Triggers HIF-1 α Causing Inflammation and Insulin Resistance in Obesity. *Cell*. 2014 June 5; 157(6): 1339–1352
3. Reference value of transcutaneous oxygen measurement in diabetic patients compared with nondiabetic patients. *J Vasc Surg* 2008;48:382-8.
4. Hyperbaric oxygen therapy improves peripheral insulin sensitivity in humans. *Diabet Med*. 2012 Aug;29(8):986-9.
5. Obstructive Sleep Apnea and Incident Diabetes. A Historical Cohort Study. *American Journal of Respiratory and Critical Care Medicine* Volume 190 Number 2, July 15, 2014
6. Sleep Apnea in Type 2 Diabetes. *Diabetes Spectrum* 2016 Feb; 29(1): 14-19.


Definitions

a) **Inflammatory response of immune system**

– When the immune system is triggered, it will release armies of immune cells into the bloodstream to fight foreign invaders such as bacteria. This is a normal response to foreign invasion. However, long term (chronic) inflammatory response of the immune system can be triggered in the absence of invasion, by substances that the body cells produce during prolonged states of stress, substances called chemokines. This continual influx of immune cells will create a state of 'martial law' in the body, causing prolonged stress to normal cells, leading to chronic diseases such as heart disease, diabetes and arthritis.

b) **Microangiopathy** – this is a complication of diabetes where the small vessels (capillaries) are damaged by high blood sugar, and they become narrowed and blocked, preventing smooth flow of blood through them.

c) **Hyperbaric oxygen chamber** – Hyperbaric Oxygen Therapy (HBOT) is a treatment that involves placing the patient inside a pressurized chamber with 100 percent oxygen. While the patient is inside the pressure chamber, it increases plasma oxygen concentration by 10 to 15 times and enables increased tissue oxygenation of the body. The oxygen concentration in plasma and the tissue oxygen levels returns rapidly to original levels within 10 minutes of the patient leaving the pressurized chamber. This is an established method of enhancing tissue oxygen for the duration that the patient is inside the pressurized chamber.



HYPOXIA IS CANCER'S BEST FRIEND

A common feature of most cancers is a low level of oxygen, called hypoxia, the severity of which varies between tumor types.¹ Cancers are hypoxic (having low level of oxygen inside the cancer tissue), the cancer cells have adapted to thrive a low oxygen environment.² This relationship between cancer and hypoxia has been known since the 1930s when Dr Otto Warburg, a German Nobel Prize winner in Medicine, discovered that cancer proliferation occurs in the absence of oxygen.³

Cancers that have lower levels of oxygen are more aggressive in behavior, with greater tendencies to invade and spread⁴, resulting in poorer outcomes for patients. Several mechanisms have been proposed by cancer researchers on how hypoxia affects cancer aggression.

In environments with low oxygen, cancer cells undergoes frequent genetic distortions (genetic mutation^a);⁵ to become aggressive cells with a ferocious ability to spread⁵. Under hypoxic conditions, cancer cells take on very primitive shapes, literally sprouting limbs that allow them to move and invade into surrounding normal tissue⁶. At the same time, they also 'give birth' to numerous immature cancer stem cells that evade detection by the immune system, hide in various parts of the body to start new colonies of tumours⁷. Non-cancer stem cells are immature cells with the ability to multiply indefinitely, to replenish tissues throughout the life of the person⁸. Cancer stem cells, unfortunately, also behave the same way to ensure the survival and recurrence of cancer in the patient's body⁷.



Professor Gregg Semenza, from Johns Hopkins Kimmel Cancer Centre puts it this way 'There are still many questions left to answer but we now know that oxygen poor environments (hypoxia), like those often found in advanced human breast cancers serve as nurseries for the birth of cancer stem cells.'⁹

According to Professor Semenza, "Chemotherapy may kill more than 99 percent of the cancer cells in a tumor but fail to kill a small population of cancer stem cells that are responsible for subsequent cancer relapse and metastasis."⁹

Cancers use hypoxia as a protective shield against radiation therapy and chemotherapy¹⁰. The more hypoxic a cancer is, the more it is resistant to treatment¹⁰. Oxygen has a substantial impact on treatment response, and hypoxia presents a serious barrier to successful radiation therapy.¹¹

Radiation therapy is ineffective against hypoxic cancer because radiation interacts with oxygen inside the cancer cell to disrupt the cell's ability to multiply¹¹. If the cancer cell is hypoxic, radiation is unable to effectively destroy it¹¹. On the other hand, having sufficient oxygen can make a big difference to how successful radiotherapy is for treating the cancer; for conventional radiation therapy, regions of a tumour with high oxygen concentration are up to threefold more responsive to treatment than regions without oxygen¹².

Failure to respond to chemotherapy is also contributed by cancer hypoxia¹³. The effective delivery of cancer drugs into hypoxic areas of the cancer is hampered by low oxygen levels and acidic conditions in the cancer tissue due to anaerobic glycolysis^{d,13}. Some chemotherapeutic drugs require oxygen to generate free radicals^c that kill cancer cells and are ineffective when the cancer is very hypoxic¹³.


Overall, hypoxia in the environment of a cancer is a critically important factor which promotes recurrence of malignancy and has a negative effect on response to most cancer treatments.

Definitions

- a) **Genetic mutation**– Mutation is a change in the cell DNA that creates slightly different versions of the same genes.
- b) **Metastasis** – Metastasis is the medical term for cancer that spreads to a different part of the body from where it started.
- c) **Free radicals** - Free radicals are atoms or groups of atoms with an odd (unpaired) number of electrons. Once formed these highly reactive radicals can start a chain reaction, that can damage cell membranes and cell DNA.
- d) **Anaerobic glycolysis** – the metabolic method used by cells to produce energy in the absence of oxygen. It produces lactic acid as a by-product.

References:

1. The role of hypoxia in cancer progression, angiogenesis, metastasis, and resistance to therapy. *Hypoxia* 2015;3:83–92
2. Hypoxia-inducible factors: mediators of cancer progression and targets for cancer therapy. *Trends Pharmacol Sci.* 2012;33(4):207–214.
3. Drivers of the Warburg phenotype. *Cancer J.* 2015 Mar-Apr;21(2):56-61
4. Molecular mechanisms and clinical applications of angiogenesis. *Nature.* 2011;473(7347):298–307.
5. Tumor hypoxia as a driving force in genetic instability. *Genome Integr.* 2013;4(1):5.
6. Complex networks orchestrate epithelial mesenchymal transitions. *Nat Rev Mol Cell Biol.* 2006;7(2):131–142
7. The hypoxic microenvironment: A determinant of cancer stem cell evolution. *Inside the Cell,* 2016, 1, 96–105
8. Stem cells for organ repair. Support or replace? *Organogenesis.* 2011 Apr-Jun; 7(2): 95.
9. Hypoxia induces the breast cancer stem cell phenotype by HIF-dependent and ALKBH5-mediated m6A-demethylation of NANOG mRNA. *PNAS* April 5, 2016. 113 (14) E2047-E2056
10. Hypoxia in prostate cancer: A powerful shield against tumour destruction? *Cancer Treatment Reviews* (2008) 34, 313– 327
11. Hall EJ: *Radiobiology for the Radiologist* (ed 3). Philadelphia, Lippincott, 1988
12. Hypoxia imaging and radiotherapy: bridging the resolution gap. *British J Radiol* 2017; 90: 2016
13. Effects of hypoxia on human cancer cell line chemosensitivity. *BMC Cancer* 2013, 13:331




FATTY LIVERS ARE HYPOXIC!

Non-alcoholic fatty liver diseases (NAFLD) is the most common long term liver disease in developed countries¹.

In this condition, the fatty changes in the liver are not due to alcohol and interestingly, a high fat diet actually reduces oxygen supply to the liver cells². Oxygen, or rather the lack of it, is a central component of the disease process leading to formation of fatty liver³. Accumulating medical evidence from the past few decades provides strong support that interruptions in oxygen supply to the liver contributes to the start and progression of fatty change³.

The liver is a highly metabolic organ, requiring a high energy supply to fuel its metabolic activities². As such, adequate oxygen supply to the liver is extremely critical for this tissue's function because without oxygen, cells produce 16 times less energy (ATP)⁴. Oxygen regulates metabolic activities in certain parts of the liver and when disease processes affect the liver, oxygen can make adjustments to how liver diseases progress⁵.



Low supply of oxygen to these parts of the liver lead to hypoxic damage of the liver cells. How well oxygenated the liver cells are prior to suffering any stress or injury can dramatically affect the outcome with better oxygenated liver cells suffering less harm⁶.

People with obstructive sleep apnea, the serious snorers, have been shown to develop fatty liver.^{7,8} The intermittent 'strangulation' occurring throughout the night in severe snorers keeps disrupting oxygen supply to the body and creates sufficient overall hypoxia to reduce supply of oxygen-rich blood to the liver and cause liver hypoxia⁹. Hypoxia then stimulates fatty change within the liver cells.

Chronic hypoxia not only propagates fatty liver, it can induce formation of fibrous tissue within the liver, leading to development of liver cirrhosis.¹⁰


References:

1. NAFLD and hepatocellular Carcinoma: how big a problem is this really? *Curr. Hepatol. Rep.* 13 (2) (2014) 113–118.
2. High fat diet induces dysregulation of hepatic oxygen gradients and mitochondrial function in vivo, *Biochem J.* 417 (1) (2009) 183–193
3. Non-alcoholic fatty liver disease, to struggle with the strangle: Oxygen availability in fatty livers. *Redox Biology* 13 (2017) 386–392
4. Oxidative Phosphorylation. D.A. Bender. *Encyclopedia of Food Sciences and Nutrition* (Second Edition), 2003
5. Oxygen: modulator of metabolic zonation and disease of the liver, *Hepatology* 31 (2) (2000) 255–260
6. Effects of hepatic zonal oxygen levels on hepatocyte stress responses, *J. Surg. Res.* 145 (1) (2008) 150–160
7. Symptoms of obstructive sleep apnea in patients with nonalcoholic fatty liver disease, *Dig. Dis. Sci.* 50 (12) (2005) 2338–2343
8. Obstructive sleep apnea is associated with fatty liver and abnormal liver enzymes: a meta-analysis, *Obes. Surg.* 23 (11) (2013) 1815–1825
9. Obstructive sleep apnea and non-alcoholic Fatty liver disease: is the liver another target? *Front. Neurol.* 3 (2012)149.
10. Hypoxia, hypoxia-inducible factors and fibrogenesis in chronic liver diseases. *Histol Histopathol.* 2014;29:33–44.

HOW IS HYPOXIA INVOLVED IN HEART ATTACK AND STROKE?

Hypoxia is the cellular weapon that kills the heart and brain during heart attacks¹ and strokes².

Heart muscle cells are very sensitive to lack of oxygen (hypoxia) because they require a constant large supply of energy to pump non-stop 24 hours a day, every day of your life. The brain is a very metabolically active organ yet it contains virtually no oxygen reserve². Brain cells are critically sensitive to hypoxia² because they are constantly transmitting electrical signals that coordinate organ function, enable movement and orchestrate other bodily functions within a living person. These activities are highly energy intensive. As we have discovered in the preceding chapter explaining why oxygen is important, cells generate 16 times more energy when well oxygenated, compared to when they try to create energy in the absence of oxygen (anaerobic glycolysis^a)³.



High functioning cells such as heart muscle cells and brain cells have high energy requirements all the time. That is why hypoxia is an undesirable state for heart and brain function.

The cardiovascular system, also known as the circulatory system, includes the heart, arteries, veins, capillaries and blood. The heart is literally the pump that moves blood through the network of blood vessels; tubes of various sizes (arteries, veins and capillaries) to reach different regions of the body.

Supplying oxygen to the body is the most essential function of the cardiovascular system. All cells in the body require oxygen to generate energy (ATP^b) in order to stay alive and perform their functions. The heart and its network of blood vessels is the transport system to ensure that oxygen is continually delivered to every cell. Although all cells require oxygen, brain cells are the most sensitive and begin to die in as little as few minutes if deprived of oxygen².



The heart pump itself uses 5 to 20 percent of the body's supply of oxygen⁴ and has its own network of blood vessels called the coronary arteries that supply blood to the heart muscles. Heart muscle cells die within 20 minutes⁴ if they do not receive enough oxygen. So hypoxia created by blocked arteries is especially lethal to heart and brain cells. According to the WHO (World Health Organization), cardiovascular disease (CVD) is the number one cause of death in the world⁵, with more people dying annually from CVDs than from any other cause. Cardiovascular disease generally refers to conditions that involve narrowed or blocked blood vessels, leading to chest pain (angina), heart attack or stroke.

What happens during a heart attack or stroke?

A heart attack is precipitated when a narrowed coronary artery suddenly becomes blocked with a blood clot and oxygen supply to the heart muscle is cut off. The heart muscle cells suffer immediate hypoxia.¹ Without oxygen, the heart muscle cells switch to anaerobic glycolysis to generate energy, making 16 times less energy (ATP) than before³. Heart muscle cells are very sensitive to low oxygen because they cannot afford to rest, they need to keep pumping blood for the rest of the body. Muscular pumping needs lots of energy (ATP). With this sudden energy crisis, the heart muscle cells become stressed and start dying. If a supply of oxygen-rich blood is not re-established within an hour, the heart muscle cells become dead and cannot be revived⁶.

If large portions of the heart muscle are dead and non-functioning, the heart pump becomes disabled. Being the main pump that pushes blood around the body, a failing heart leads to poor oxygen supply to the entire body. This is a state called heart failure. If the heart pump stops because of extensive damage, the person will die.

A stroke is simply the same event described above, happening to an artery supplying blood to the brain. Hypoxia in brain cells is a very serious problem because brain cells die even faster than heart muscle cells when hypoxia sets in - dying within a few minutes from the time oxygen supply is cut off². This leads to rapid cell death and permanent damage to the affected parts of the brain within minutes². Depending on which region of the brain the affected artery supplies to, a stroke can result in a range of different neurological problems such as loss of ability to move a limb, loss of ability to speak, loss of sight or even loss of consciousness (coma). Damage in critical parts of the brain from a stroke can lead to death.



Definitions

- a) **Anaerobic glycolysis** - Anaerobic glycolysis is a method used by cells to produce ATP by transforming glucose into lactate when limited amounts of oxygen (O₂) are available. Compared to OXPHOS, anaerobic glycolysis creates 16 times less energy.
- b) **ATP** - Adenosine triphosphate, an energy-carrying molecule found in the cells of all living things. ATP captures chemical energy obtained from the breakdown of food molecules and releases it to fuel all cellular processes needed for life. It is the source of energy that keeps everything going.

References:

1. Cellular Pathways of Death and Survival in Acute Myocardial Infarction. J Clin Exp Cardiol 2012, S:6
2. Hypoxia, Ischemic Stroke, and Memory Decits: Prospects for Therapy. IUBMB Life, 48: 373-378, 1999
3. Oxidative Phosphorylation. D.A. Bender. Encyclopedia of Food Sciences and Nutrition (Second Edition), 2003
4. 2000-2018 The Cleveland Clinic Foundation, Centre for Continuing Education.
5. WHO statistics 2015
6. Time to Treatment in Primary Percutaneous Coronary Intervention. N Engl J Med 2007;357:1631-8



WOUND HEALING SUFFERS WHEN THERE IS HYPOXIA

Oxygen has a massive effect on the wound healing process, and one of the biggest factors that can inhibit the body's ability to heal a wound is low oxygen levels (hypoxia) in the affected area¹.

While the exact way oxygen works in the wound healing process is more complex than understood, it is widely recognized that oxygen is needed in nearly every stage of the wound healing pathway².

When the body acquires a wound, from surgery or trauma, the break in the body's integrity makes the wound vulnerable and it triggers a response to greater bacterial defense, cell proliferation, collagen synthesis and regeneration of blood vessels and nerves, among other reparative activities³.

All these are high energy activities. As reported in the British Journal of Dermatology³, oxygen's main function in wound healing lies in its capacity to effectively produce energy. In order for cells to fight infections, properly multiply and get organized into different skin cells to fill up the wound, they must have a sufficient amount of energy¹.

Like all functions involved in human biology, oxygen is essential for cells to generate energy (ATP^a) effectively, using a process called oxidative phosphorylation^b.

When an area of the body does not receive an adequate amount of oxygen, a condition known as hypoxia, it can slow and even halt the healing process – resulting in chronic wounds^c.

The limitation of oxygen delivery to the wound is often due to many factors; however, the end result is always non-healing, chronic wounds or ulcers. Chronic leg and foot ulcers occur in many adults with vascular disease or diabetes and are attributed to poor delivery of oxygen rich blood through blocked blood vessels, prolonged pressure, or abnormal nerve function.⁴

These ulcers last on average 12 to 13 months, can relapse in up to 60% to 70% of patients, can lead to loss of function, amputation and decreased quality of life.⁴ These wounds become infected because there is insufficient oxygen to mount an effective antibacterial effect and ultimately, the poor oxygenation of the skin over a prolonged period leads to gangrene (death of the tissue) of the affected area and amputation of the leg⁵.



Doctors currently use hyperbaric oxygen therapy (HBOT)^d, an oxygenation method to correct hypoxia in poorly healing wounds⁶. HBOT is usually effective to heal these chronic wounds but many patients are unable to tolerate the side effects such as pressure induced ear damage (middle ear barotrauma^e)⁷, dental pains, difficulty with visual focus⁸, risks of cataracts⁹, and increased blood pressure¹⁰. Middle ear barotrauma is one of the most common side effects of HBOT. Patients experience problems with ear equalization, a feeling of pressure, ear pain, and discomfort during the time they ramp up the pressure in the chamber. As such, this current method of overcoming wound hypoxia is not suitable for every patient.

Definitions

a) **ATP** – Adenosine triphosphate, an energy-carrying molecule found in the cells of all living things. ATP captures chemical energy obtained from the breakdown of food molecules and releases it to fuel all cellular processes needed for life. It is the source of energy that keeps everything going.

b) **OXPHOS** - Oxidative Phosphorylation is process whereby a cell uses oxygen to generate large amounts of energy in the form of ATP. This process, which takes place in mitochondria, is the major source of ATP in aerobic organisms.

c) **Chronic wounds** - Chronic ulcers are those that do not progress through the healing process in a timely manner and typically last 12 to 13 months.

d) **HBOT** - Hyperbaric oxygen therapy (HBOT) an adjunct therapy to increase tissue oxygen with the use of 100% oxygen at air pressures greater than atmospheric pressure.

e) **Middle Ear Barotrauma** – a common side effect of the high pressure–high oxygen environment during HBOT, leading to sensation of ear pain, ear canal swelling and rupture of eardrum with resulting deafness.

References:

1. Wound Healing Essentials: Let There Be Oxygen. *Wound Repair Regen.* 2009; 17(1): 1–18.
2. Wounds: an overview of the role of oxygen. *Antioxid Redox Signaling.* 2007;9(8):1183-1192.
3. Oxygen in acute and chronic wound healing. *British Journal of Dermatology* Volume163, Issue 2 August 2010. Pages 257-268
4. Evidence-based management of common chronic lower extremity ulcers. *Dermatol Ther* 2013;26:187–196
5. Chronic Wound Healing: A Review of Current Management and Treatments. *Adv Ther* (2017) 34:599–610
6. Hyperbaric oxygen—its mechanism and efficacy. *Plast Reconstr Surg* 2011;127(S1):131S–141S
7. Hyperbaric Oxygen Therapy: Side Effects Defined and Quantified. *Adv Wound Care* (New Rochelle). 2017 Jun 1; 6(6): 210–224.
8. Ocular complications in hyperbaric oxygen therapy. In: Neuman TS, editor, Thom SR, editor. eds. *Physiology and Medicine of Hyperbaric Oxygen Therapy.* Philadelphia, PA: Saunders Elsevier, 2008:565–572
9. Nuclear cataract and myopia during hyperbaric oxygen therapy. *B J Ophthalmol* 1984;68:113–117
10. Influences of hyperbaric oxygen on blood pressure, heart rate and blood glucose levels in patients with diabetes mellitus and hypertension. *Arch Med Res* 2006;37:991–997

INTRODUCING ELO OXYGEN-RICH DRINKING WATER

ELO Drinking Water is an oxygen-rich water with a refreshing, smooth taste that nourishes the body from within and enhances overall well-being. The revolutionary technology behind ELO Drinking Water allows a high level of oxygen to exist in a unique, stable and bound form that promotes quick absorption by the body, restoring its natural balance and enhancing overall well-being.

Properties of ELO Drinking Water:

- High oxygen content
- A pH of between 7.1 and 7.5
- Natural minerals
- Refreshing smooth taste
- Preservative-free
- Pure and additive-free

PURCHASE INFORMATION

ELO Oxygen-Rich Drinking Water is sold at ELO Lab, City Square Mall and ELO Water Therapy Centre, Belvedere and available for purchase online at the following websites. Delivery can be arranged upon request. Charges may apply.

www.elowatershop.com

